

ACBS ✓ GMS PBS

Description

Food for special medical purposes.

MMA/PA explore¹¹⁵ is an unflavoured powdered methionine, threonine and valine free*, low isoleucine protein substitute containing essential and non-essential amino acids, carbohydrate, sugar, vitamins, minerals, trace elements, arachidonic acid (ARA) and docosahexaenoic acid (DHA).

Indications

MMA/PA explore5 is for the dietary management of Methylmalonic Acidaemia / Propionic Acidaemia (MMA/PA) from 6 months to 5 years of age.

Dosage and Administration


To be determined by the clinician or dietitian and is dependent on the age, body weight and medical condition of the patient.

MMA/PA explore5 (12.5g) = 5g PE

- 1. The daily protein substitute requirement is given as MMA/PA explore5.
- 2. Additional protein is provided from exchanges of natural protein, to provide restricted amino acids in controlled amounts, which is essential for growth and development.
- 3. Foods high in carbohydrate and fat, but low in protein, should be provided to supply the remainder of the energy.

Preparation Guidelines

MMA/PA explore5 is taken as a semi-solid spoonable consistency:



- 1. Add 12.5ml of cold water into a beaker.
- 2. Empty the contents of the sachet into the beaker.
- 3. Secure the lid and shake well for 10 seconds.
- $4.\,Leave$ to stand for 2 minutes, allowing a smooth semi-solid consistency to form. Consume with a spoon.

Water or permitted drinks should be taken after MMA/PA explore5.

The product should be prepared immediately prior to feeding and any remaining product must be discarded if not used within 1hr.

IMPORTANT NOTICE

Must be used under strict medical supervision.

Not suitable for use as a sole source of nutrition.

Suitable from 6 months to 5 years of age.

Must only be consumed by children with proven Methylmalonic Acidaemia/ Propionic Acidaemia (MMA/PA).

Diet must be supplemented with natural protein, water and other nutrients in prescribed quantities to supply methionine, threonine, valine, isoleucine, fluid and general nutritional requirements.

For enteral use only.

Nutritional	MMA/PA	explore:	5		MMA/PA	A explore5
Information	per 100g	per 12.5g			per 100g	per 12.5g
Energy kJ	1450	181	Vitamin B12	μд	3.7	0.46
kcal	342	43	Pantothenic A	cid mg	7.3	0.91
Fat g	1.5	0.2	Biotin	μд	23	2.9
of which saturates g	0.7	0.09	Sodium	mg	195	24
ARA mg	280	35		mmol	8.4	1.0
DHA mg	140	18	Chloride	mg	500	63
Carbohydrate g	42	5.3		mmol	14	1.8
of which sugars g	28	3.5	Potassium	mg	500	63
Protein equivalent g	40	5.0		mmol	12	1.6
L-Methionine* g	-	-	Calcium	mg	1100	140
L-Threonine* g	-	-		mmol	28	3.4
L-Valine* g	-	-	Phosphorus	mg	730	91
L-Isoleucine g	0.13	0.02		mmol	23	2.9
Salt g	0.49	0.06	Magnesium	mg	135	17
Vitamins and minera	als			mmol	5.5	0.69
Vitamin A (RE) μg	530	66	Iron	mg	16.0	2.0
Vitamin D μg	27	3.4	Zinc	mg	11.0	1.4
Vitamin E (αTE) mg	9.7	1.2	Copper	mg	0.78	0.10
Vitamin K μg	27	3.4	lodine	μд	150	19
Vitamin C mg	90	11	Selenium	μд	31	3.9
Thiamin mg	1.0	0.12	Manganese	mg	0.25	0.03
Riboflavin mg	1.5	0.19	Chromium	μд	30	3.8
Vitamin B6 mg	1.1	0.14	Molybdenum	μд	40	5.0
Niacin mg (mg NE)	6.9 (26)	0.86 (3.3)	Others			
Folic Acid µg	230	29	Choline	mg	450	56

Nutritional		MMA/PA explore5				MMA/PA explore5		
Information		per 100g	per 12.5g			per 100g	per 12.5g	
Amino acids				L-Lysine	g	4.52	0.57	
L-Alanine	g	4.12	0.52	L-Phenylalanine	g	2.79	0.35	
L-Arginine	g	4.23	0.53	L-Proline	g	2.04	0.26	
L-Aspartic Acid	g	7.35	0.92	L-Serine	g	3.05	0.38	
L-Cystine	g	1.79	0.22	L-Tryptophan	g	1.15	0.14	
L-Glutamine	g	5.01	0.63	L-Tyrosine	g	2.79	0.35	
Glycine	g	1.96	0.25	L-Carnitine	mg	44	5.5	
L-Histidine	g	2.79	0.35	Taurine	mg	88	11	
L-Leucine	g	5.81	0.73					

^{*} No added methionine, threonine or valine. These may be present in trace amounts from other ingredients (<10mg/100g powder, <4mg/serving).

Allergen Declaration

Contains Fish (Tuna). Not suitable for individuals with milk allergy.

Pack Size/Weight



30 x 12.5g **e** sachets = 375g

Flavours

Unflavoured.

Storage

Store in a cool, dry place. Sachets are designed for single use. Any unused powder should be tightly sealed and used within 24 hours.



