

# PEPTAMEN® JUNIOR 0.6

## DESCRIPTION

Peptamen® Junior 0.6 is a 0.6 kcal/ml peptide-based enteral tube feed for the dietary management of disease related malnutrition in patients with malabsorption and/or maldigestion who have reduced energy requirements. It is a partially hydrolysed, 100% whey protein formula with omega-3 fatty acids, fibre and 35.6% of the fat as MCT. Peptamen® Junior 0.6 is a nutritionally complete Food for Special Medical Purposes for use under medical supervision.

## PRESENTATION

500ml SmartFlex® collapsible semi-rigid bottles, 12 x 500ml per case.

## INDICATIONS

ACBS approved, prescribable on FP10 (GP10 in Scotland). Indicated as a sole source of nutrition for children aged 1 year onwards for the dietary management of disease related malnutrition in patients with malabsorption and/or maldigestion who have reduced energy requirements.

## INGREDIENTS

Water, maltodextrin, peptides from partially hydrolysed whey protein (milk), soybean oil, starch, sucrose, MCT oil, fibres (partially hydrolysed guar gum, pea fibre), minerals (potassium citrate, calcium phosphate, magnesium chloride, sodium chloride, calcium citrate, sodium phosphate, magnesium oxide, ferrous lactate, zinc sulphate, copper gluconate, manganese sulphate, sodium fluoride, potassium iodide, chromium chloride, sodium selenite, sodium molybdate), emulsifiers (E472c, E471), fish oil, choline bitartrate, vitamins (C, niacin, E, pantothenic acid, B<sub>6</sub>, riboflavin, thiamin, A, folic acid, biotin, K, D, B<sub>12</sub>), stabiliser (E407), taurine, inositol, L-carnitine.

## ADMINISTRATION AND DOSAGE

Intended for children with identified reduced energy needs. Recommended volumes should be based on a thorough assessment and continuous evaluation of the needs of the child.

## DIRECTIONS FOR USE AND STORAGE

Shake well before use. Once spiked, use within 24 hours. Recap unused portion, refrigerate and use within 24 hours. If poured out of the pack, cover tightly and use within 6 hours. In order to maintain a constant flow, the use of a pump is recommended. Unopened, store in a cool dry place.

## CONTRAINDICATIONS

Not suitable for patients with a cows' milk allergy.

## PRECAUTIONS

For enteral use only. Suitable for children above 1 year of age. Do not add food or drugs to the feed.



# NUTRITION INFORMATION

Typical values	Per 100ml	Per 500ml
Energy kJ/kcal	254 / 60	1270 / 300
Fat (37% kcal) g	2.5	12.5
of which saturates g	1.3	6.5
- MCT g	0.89	4.4
of which monounsaturates g	0.26	1.3
of which polyunsaturates g	0.64	3.2
- Omega-3 mg	86	430
Carbohydrate (45% kcal) g	6.8	34
of which sugars g	1.5	7.5
- Lactose g	< 0.2	
Fibre (3% kcal) g	0.80	4.0
Protein (15% kcal) g	2.3	11.5
Salt g	0.13	0.65
<b>Vitamins</b>		
A µg	50	250
D µg	1.6	8.0
E mg	0.9	4.5
K µg	4.2	21
C mg	6.0	30
Thiamin mg	0.08	0.40
Riboflavin mg	0.12	0.60
Niacin mg (mg NE)	1.04 (1.7)	5.2 (8.5)
B6 mg	0.10	0.50
Folic acid µg	14	70
B12 µg	0.24	1.2
Biotin µg	2.7	13.5
Pantothenic acid mg	0.44	2.2
<b>Minerals</b>		
Sodium mg/mmol	52/2.3	260/11
Potassium mg/mmol	126/3.2	630/16
Chloride mg/mmol	77/2.2	385/11
Calcium mg/mmol	68/1.7	340/8.5
Phosphorus mg	34	170
Phosphate mmol	1.1	5.5
Magnesium mg/mmol	20/0.83	100/4.2
Iron mg	0.86	4.3
Zinc mg	0.61	3.0
Copper mg	0.07	0.35
Manganese mg	0.14	0.70
Fluoride mg	0.070	0.35
Selenium µg	3.0	15
Chromium µg	2.4	12
Molybdenum µg	5.0	25
Iodine µg	11	55
<b>Other Nutrients</b>		
Choline mg	25	125
Taurine mg	7.5	37.5
L- Carnitine mg	4.0	20
Osmolarity mOsm/l	206	
Water content per 100ml:	90 g	

NE = Niacin Equivalent