Description

Oral Impact® is a powdered enteral sip feed that provides 1.0 kcal/ml when reconstituted with water. Nutritionally complete Food for Special Medical Purposes with omega-3 fatty acids, arginine, nucleotides and soluble fibre.

POWDER WITH THE POWDER

Presentation

5 sachets x 74g per box. Available in Tropical, Coffee and Citrus flavour.

Indications

ACBS approved, prescribable on FP10 (GP10 in Scotland). Indicated for the preoperative nutritional support of patients who are malnourished or at risk of becoming malnourished.

Ingredients - Tropical Flavour*

Sucrose, whey protein (milk), maltodextrin, L-arginine, fish oil, fibre (guar gum), minerals (potassium citrate, sodium chloride, calcium phosphate, magnesium citrate, potassium dihydrogen phosphate, ferrous sulphate, zinc sulphate, copper gluconate, manganese sulphate, chromium chloride, potassium iodide, sodium selenite, sodium molybdate, sodium fluoride), medium chain triglycerides, corn oil, acidity regulator (E330), yeast extract rich in ribonucleotides (RNA), stabiliser (E415), choline bitartrate, emulsifier (soy lecithin), vitamins (C, E, pantothenic acid, niacin, B6, B1, B2, A, folic acid, D, K, biotin), flavourings, antioxidants (E304, E306, E 307, contains soy lecithin), colours (beetroot red, beta carotene).

Administration and dosage

Dissolve contents of one sachet in 250 ml of cool boiled water. Use shaker, whisk or fork to mix. Consume immediately. The recommended pre-operative dose is: 500-1000ml (500-1000 kcal) per day + usual diet for 5-7 days prior to surgery, dependent on nutritional requirements (Waitzberg DL et al, 2006). The recommended post-operative dose is: 1200-1500ml (1200-1500 kcal) per day for 5-7 days post-surgery, dependent on nutritional requirements.

Shelf life and storage

12 months. Should be stored unopened in a cool, dry place. Store between 4°C – 25°C away from light and humidity.

Contraindications

Suitable for children above 3 years of age. For oral use only. Do not administer to severely septic patients. Do not mix with medication and food.

Precautions

Must be used under medical supervision.

Nutrition information - Tropical flavour*

| Tunical values | Dox 100# | Dox 7/a |
|----------------------------------|----------|-----------|
| Typical values | Per 100g | Per 74g |
| | | (Tropical |
| | | flavour*) |
| General | | |
| Energy kJ/kcal | 1760/418 | 1304/309 |
| Fat (22% kcal) g | 10 | 7 |
| of which saturates g | 5.1 | 4 |
| of which monounsaturates g | 2.1 | 1.5 |
| of which polyunsaturates g | 3.2 | 2.3 |
| Carbohydrates (54% kcal) g | 56 | 41.4 |
| of which sugars g | 30 | 22 |
| of which lactose g | 1.2 | 0.89 |
| Fibre (2% kcal) g | 4.1 | 3 |
| Protein (23% kcal) g | 24 | 18 |
| Salt (= Na(g) x 2.5) | 1.1 | 0.8 |
| Vitamins | | |
| A μg RE | 520 | 385 |
| Dμg | 4.2 | 3.1 |
| К µg | 27 | 20 |
| C mg | 105 | 78 |
| B1 (Thiamin) mg | 0.49 | 0.36 |
| B2 (Riboflavin) mg | 0.8 | 0.59 |
| B6 mg | 0.7 | 0.52 |
| Niacin mg/mg NF | 1.7/6.5 | 1.3/4.8 |
| Niacin mg/mg NE Folic acid µg | 140 | 104 |
| B12 μg | 2.5 | 1.8 |
| Pantothenic acid mg | 4 | 3 |
| Biotin µg | 20 | 15 |
| E mg α-TE | 12.2 | 9 |
| Minerals | | , |
| Sodium mg/mmol | 434/19 | 321/14 |
| Chloride mg/mmol | 608/17 | 450/13 |
| Potassium mg/mmol | 543/13.9 | 402/10 |
| Calcium mg/mmol | 324/8.1 | 240/6.0 |
| Phosphorus mg/mmol | 292/9.4 | 216/7.0 |
| Magnesium mg/mmol | 70/2.9 | 52/2.2 |
| Iron mg | 4.9 | 3.6 |
| Zinc mg | 5 | 3.7 |
| Copper mg | 0.69 | 0.51 |
| lodine µg | 61 | 45 |
| Selenium µg | 25 | 18 |
| Manganese µg | 810 | 599 |
| Chromium µg | 41 | 30 |
| Molybdenum µg | 50 | 37 |
| Fluoride µg | 540 | 400 |
| Other nutrients | | |
| Choline mg | 165 | 122 |
| L-arginine g | 5.1 | 3.8 |
| Omega-3 g | 1.3 | 0.96 |
| MCT g | 3 | 2.2 |
| Nucleotides g | 0.6 | 0.45 |
| Osmolarity mOsm/l | 520 | 0.43 |
| Osmolality mOsm/kg H20 | 620 | |
| OSITIOTALITY IIIOSIII/ Kg HZU | 020 | |

^{*} See individual packs for other flavours